

Table 2. Estimated distribution of persons with and persons without a diagnosis of AIDS within 12 months of diagnosis of HIV infection, by selected characteristics, 2001—30 areas with confidential name-based HIV infection reporting

	AIDS diagnoses				Total	
	≥12 Months after diagnosis of HIV infection		<12 Months after diagnosis of HIV infection			
	No.	(%) ^a	No.	(%) ^a	No.	(%) ^a
Age at diagnosis (yrs)						
<13	160	77	47	23	206	100
13—14	25	75	8	25	33	100
15—24	2,427	83	500	17	2,926	100
25—34	4,757	66	2,464	34	7,221	100
35—44	5,175	57	3,942	43	9,118	100
45—54	2,210	50	2,198	50	4,408	100
55—64	577	44	726	56	1,303	100
≥65	189	44	238	56	427	100
Race/ethnicity						
White, not Hispanic	4,817	60	3,222	40	8,039	100
Black, not Hispanic	8,627	61	5,463	39	14,090	100
Hispanic	1,719	57	1,293	43	3,012	100
Asian/Pacific Islander	82	58	58	42	140	100
American Indian/Alaska Native	74	52	68	48	142	100
Exposure category						
Male adult or adolescent						
Male-to-male sexual contact	6,461	59	4,512	41	10,973	100
Injection drug use	1,419	52	1,296	48	2,715	100
Male-to-male sexual contact and injection drug use	554	58	394	42	948	100
Heterosexual contact	1,858	58	1,333	42	3,191	100
Other ^b	81	48	87	52	167	100
Subtotal	10,373	58	7,621	42	17,994	100
Female adult or adolescent						
Injection drug use	948	64	533	36	1,481	100
Heterosexual contact	3,974	68	1,852	32	5,826	100
Other ^b	65	48	70	52	136	100
Subtotal	4,987	67	2,455	33	7,442	100
Child(<13 yrs)						
Perinatal	148	77	44	23	192	100
Other ^c	12	83	2	17	14	100
Subtotal	160	77	47	23	206	100
Total^d	15,520	61	10,123	39	25,642	100

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk. The estimates have not been adjusted for incomplete reporting.

Data include persons in whom AIDS has developed and persons whose first diagnosis of HIV infection and the diagnosis of AIDS were made at the same time.

Data exclude 1 person whose month of HIV infection diagnosis is unknown.

Since 1998, the following 30 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Percentages represent proportions of the total number of diagnoses of HIV/AIDS made during 2001 for the corresponding group (see row entries).

^b Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk not reported or not identified.

^d Includes 220 persons of unknown or multiple race. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.